

Moisture index during the last two centuries inferred from tree growth in the western Himalaya, India

Somaru Ram* and H. P. Borgaonkar

Indian Institute of Tropical Meteorology, Pune 411 008, India

In the present study, a regional tree-ring chronology prepared from two species (*Picea smithiana* and *Cedrus deodara*) over the western Himalaya has been used in relation to climate fluctuations. This analysis shows that moisture index and rainfall during February to May have significant positive relationship, whereas temperature and heat index show a negative relationship with tree growth. However, moisture and heat indices show greater impact on tree growth than rainfall and temperature. The strong association of tree-ring chronology with moisture demonstrates that tree rings are much more sensitive to the availability of moisture at the root zone, which enabled us to extend our analysis back to AD 1789; in the present reconstruction, moisture deficiency for two consecutive years was noticed during 1846–1847, 1908–1909, 1921–1922, 1931–1932, 1947–1948 and 1966–1967.

Keywords: Moisture and heat indices, rainfall, temperature, tree rings.

THE western Himalaya has diverse climate (moist, moist subhumid, dry subhumid and semiarid) due to the geographical setting, high mountain ranges, variation in topography and orography. There is a large-scale variation in precipitation at short distances with changing mountain ranges and elevation. In the case of temperature, it is found to have a strong association spatially¹. However, to understand the long-term variations in precipitation and temperature over the region, proxy records of several tree ring-width samples of conifers from the Himalaya have been performed in relation to climate variability/change^{2–7}. They have shown good potential of Himalayan tree ring data for dendroclimatic analysis. However, all these studies are based on rainfall and temperature only. Reconstruction of moisture index (MI), which is a function of rainfall and temperature⁸, is not yet available in a longer perspective over the region. Therefore, in the present study, we have reconstructed MI for the last two centuries. Such climatic records may help understand the past behaviour of availability of moisture at the root zone of trees.

In this context, ring-width data of *Picea smithiana* and *Cedrus deodara* from two sites (Dhanolti and Jageswar) in the western Himalaya have been used to study tree growth–climate relationship^{3,4} (Figure 1). Site informa-

tion, number of core samples per tree and their standardization techniques have already been discussed in a previous study⁴; consistently strong correlation was observed during the chronology length⁴. However, to check the spatial coherence in chronology, inter-correlations were performed for the period 1838–1990, based on a 50-year window with overlapping of 25 years. Correlation coefficients (CCs) ranged from 0.28 to 0.61, and were significant at 5% level, indicating common forcing, i.e. climate. Such strong correlations enabled us to prepare a regional chronology by merging tree core samples of the two sites to develop a robust climate reconstruction (Figure 1). A total of 25 core samples were considered in the present study and their ring-width measurement and crossdating were checked using COFECHA software⁹.

All ring-width series were detrended using ARSTAN¹⁰. To obtain the common signal among the samples, double detrending process was adopted for all the series. First, ring-width data were standardized by a linear or negative exponential method to remove biological growth¹¹. The resulting indices were detrended once again by 35-year cubic smoothing spline for reducing the remaining trends¹¹. The obtained indices were prewhitened using autoregressive model based on the Akai information criterion to minimize the persistent low-frequency signal¹¹, and combined these indices across all the series to prepare a mean chronology¹².

Tree-ring data like mean sensitivity (0.21), variance (0.31), expressed population signal (0.87) and signal-to-noise ratio (6.5) showed the greater impact of climate on the tree rings. The threshold value (0.85) of the expressed population signal (EPS) was found to be above that reported since AD 1788 (figure not shown), which indicated reliability and stability of chronology for dendroclimatic analysis^{3,12}. However, the noisy data associated with poor replication of tree core samples were discarded and chronology was truncated to AD 1788 based on the performance by EPS^{3,12}.




Figure 1. Map showing the study area. $\Delta 1$ and $\Delta 2$, Tree-ring width sites; \bullet , Grid point climatic data.

*For correspondence. (e-mail: somaru@tropmet.res.in)

Table 1. Statistical tests performed during the calibration and verification periods for reconstruction of moisture index

Period	Calibration			Verification			
	F	R	V	Period	r	Pmt	RE
1901–1945	39.2**	0.69**	47.6%	1946–1990	0.50**	6.26**	0.47
1946–1990	11.6**	0.48**	23%	1901–1945	0.70**	3.97**	0.43
1901–1990	47.4**	0.60**	36%				

**Significant at 0.1%.

R, Multiple correlation coefficients; V, Variance explained; r, Correlation coefficients between actual and reconstructed moisture index; Pmt, Product mean test; RE, Reduction of error¹⁹.




Figure 2. Correlations of tree-ring chronology with heat index HI (—●—), Temperature (TM) (—), rainfall (PPT) (—■—) and moisture index (MI) (—◆—). Dashed lines are significant at 5% level.




Figure 3. a, Variations in actual (solid) and reconstructed MI (dashed). CC, Correlation coefficient. b, Reconstructed MI during 1789–1990. The smoothed line shows 30-year cubic spline fit.

To analyse the relationship between chronology and climate data, the grid boxes (30.75°N, 78.25°E; 30.25°N, 78.25°E; 30.25°N, 78.75°E; 29.25°N, 78.75°E; 29.75°N, 78.75°E; 29.25°N, 79.25°E; 29.75°N, 79.25°E) of rainfall

and temperature have been taken from CRU¹³ (Figure 1). A regional series for precipitation and temperature was prepared for the period 1901–2002. In addition, MI and heat index (HI) were computed in relation to trees growth variations^{8,14}.

The correlation analysis between residual chronology and HI, temperature (Tm), rainfall (PPT) and MI was performed on a monthly basis (Figure 2). In Figure 2, the dashed horizontal lines are significant at 5% level. The significant positive association between a chronology and MI and PPT was observed during current year February, March, April and May. For Tm and HI, chronology showed significant negative relationships during November and December of the previous year, and the current year March, April and May.

Based on relationship, an analysis has been made to know the impact of seasonal climate on growth variations of trees during February to May. CCs between a chronology and HI, Tm, PPT and MI were –0.53, –0.49, 0.57 and 0.60 respectively, showing significance at 0.01% level. The analysis showed that seasonal climate has greater impact on tree growth variations than a single climatic parameter^{15,16}. However, the relationship of tree growth with HI and MI was found stronger than that with temperature and rainfall. The results show that MI, which describes soil moisture availability at the root zone of the trees, plays a significant role in early wood formation¹⁷, whereas increased HI is not found conducive to tree growth due to high potential evapotranspiration¹⁶.

However, MI which showed the highest CC with tree growth was estimated to the past two centuries based on chronology of tree rings. In the regression model, tree-ring chronology with lead/lag of one year, i.e., t_{t+1} , t_0 and t_{t-1} was selected as predictor and MI as predictand¹⁸. MI was split in two sub-periods, i.e. 1901–1945 and 1946–1990, so that statistical tests could be compared against each sub-period. The first sub-period model accounted for 47.6% of the variance, whereas the calibration model of the second sub-period accounted for 23% of the variance (Table 1), weaker than the first calibration model, but showed the highly significant correlation (r), Pmt, and RE (Table 1). These results indicate reliability in tree growth–climate relationship¹⁹. Based on the various statistical tests, a calibration model for the full period

(1901–1990) was developed to extract the maximum low-frequency variation in reconstruction (Table 1). The reconstruction showed good year-to-year agreement and was highly correlated with actual data (Figure 3 a).

Figure 3 b shows the reconstructed MI back to AD 1789. Smoothed line in the figure shows cubic smoothing spline fit of 30 years. During the years 1798, 1800, 1806, 1814, 1822, 1825–1827, 1850, 1859, 1863, 1865, 1868, 1871, 1875, 1882–1883, 1886, 1889, 1893, 1905, 1912–1914, 1917, 1925–1928, 1933, 1936–1937, 1940, 1946, 1949, 1954, 1957, 1963, 1971, 1979 and 1981, high MI values (more than mean \pm 1 SD) were noticed in the present reconstruction. It shows that the moist climate (humid) during spring season plays an important role in trees growth, whereas moist subhumid climate (less than mean-1std) during 1790, 1810, 1813, 1816, 1837, 1846–1847, 1860, 1869, 1873, 1876, 1879, 1884, 1887, 1890, 1892, 1896, 1908–1909, 1921–1922, 1931–1932, 1934, 1939, 1941, 1947–1948, 1953, 1958, 1966–1967, 1972, 1974 and 1985 might suppress trees growth¹⁶.

To verify the present reconstruction, we have compared it with an earlier reconstruction by Yadav *et al.*²⁰, who also reported drought year in 1920–1924, 1812–1816, 1965–1968 and pluvial phase in 1791–1792, 1911–1915. Cook *et al.*²¹ and Ram²² also showed low soil moisture availability during 1848–1849, 1887–1897, 1933–1949 and 1965–1973, which is similar to our reconstruction. Overall, the present reconstruction shows somewhat similar patterns with previous reconstructions. However, some differences exist between the present and previous reconstructions, which could be due to geographical settings of the western Himalaya, or usage of a new climatic parameter, which is different from the previous reconstructions.

Moisture deficiency noticed during two consecutive years, i.e. 1846–1847, 1908–1909, 1921–1922, 1931–1932, 1947–1948 and 1966–1967, motivates us to prepare a longer chronology with good replication of samples, so that MI and HI could possibly be reconstructed even to further back in time. Monitoring the historical records of climate for the past several centuries over the region may be useful for a better understanding of past extreme events (driest and wettest). This type of study is useful in forest management, water resources and for sustainable development over the region.

1. Yadav, R. R., Park, W.-K. and Bhattacharyya, A., Spring temperature variations in western Himalaya, India, as reconstructed from tree-rings: AD 1390–1987. *Holocene*, 1999, **9**, 85–90.
2. Hughes, M. K., Dendroclimatic evidence from western Himalaya. In *Climate: Since AD 1500* (eds Bradley, R. S. and Jones, P. D.), Routledge, London, 1992, pp. 415–431.
3. Pant, G. B., Borgaonkar, H. P. and Rupa Kumar, K., Climatic signals from tree-rings: a dendroclimatic investigation of Himalayan spruce (*Picea smithiana*). *Himalayan Geol.*, 1998, **19**(2), 65–73.
4. Borgaonkar, H. P., Pant, G. B. and Rupa Kumar, K., Tree-ring chronologies from western Himalaya and their dendroclimatic potential. *IAWA*, 1999, **20**(3), 295–309.

5. Yadav, R. R. and Singh, J., Tree-ring based spring temperature patterns over the past four centuries in western Himalaya. *Quaternary Res.*, 2002, **57**, 299–305.
6. Singh, J. and Yadav, R. R., Spring precipitation variations over the western Himalaya, India, since AD 1731 as deduced from tree-rings. *J. Geophys. Res.*, 2005, **110**; doi:10.1029/2004JD004855.
7. Singh, J., Yadav, R. R. and Wilmking, M., A 694-year tree-ring based rainfall reconstruction from Himachal Pradesh, India. *Climate Dyn.*, 2009, doi:10.1007/s00382-09-0528-5.
8. Ram, S., Borgaonkar, H. P. and Sikder, A. B., Tree-ring analysis of teak in Central India and its relationship with rainfall and moisture index. *J. Earth Syst. Sci.*, 2008, **117**(5), 637–645.
9. Holmes, R. L., Computer assisted quality control in tree-ring dating and measurement. *Tree-Ring Bull.*, 1983, **44**, 69–75.
10. Cook, E. R., A time series analysis approach to tree-ring standardization; Dissertation, University of Arizona, Tucson, Arizona, USA, 1985.
11. Cook, E. R. and Kairiukstis, L. A. (eds), *Method of Dendrochronology: Application in Environmental Sciences*, Kluwer Academic, Dordrecht, The Netherlands, 1990, p. 394.
12. Wigley, T. M. L., Briffa, K. R. and Jones, P. D., On the average value of correlated time series, with applications in dendroclimatology and hydrometeorology. *J. Climate Appl. Meteorol.*, 1984, **23**, 201–213.
13. Mitchell, T. D. and Jones, P. D., An improved method of constructing a database of monthly climate observations and associated high resolution grids. *Int. J. Climatol.*, 2005, **25**, 693–712.
14. Thornthwaite, C. W., An approach towards a rational classification of climate. *Geogr. Rev.*, 1948, **38**, 55–94.
15. Cook, E. R., Meko, D. M., Stahle, D. W. and Cleaveland, M. K., Drought reconstructions for the continental United States. *J. Climate*, 1999, **12**, 1145–1162.
16. Ram, S. and Borgaonkar, H. P., Tree-ring analysis over the western Himalaya and its long-term association with vapor pressure and potential evapotranspiration. *Dendrochronologia*, 2014, **32**, 32–38.
17. Yadav, R. R., Long-term hydroclimatic variability in monsoon shadow zone of western Himalaya, India. *Climate Dyn.*, 2010, doi:10.1007/s00382-010-0800-8.
18. Cook, E. R., Briffa, K. and Jones, P. D., Spatial regression methods in dendroclimatology: a review and comparison of two techniques. *Int. J. Climatol.*, 1994, **14**, 379–402; doi:10.1002/joc.3370140404.
19. Fritts, *Tree-Ring and Climate*, Academic Press, London, 1976.
20. Yadav, R. R., Misra, K. G., Yadav, A. K., Kotlia, B. S. and Misra, S., Tree-ring footprint of drought variability in last 300 years over Kumaun Himalaya, India and its association with crop productivity. *Quaternary Sci. Rev.*, 2015, **117**, 113–123.
21. Cook, E. R., Anchukaitis, K. J., Buckley, B. M., D'Arrigo, R. D., Jacoby, G. C. and Wright, W. E., Asian monsoon failure and megadrought during the last millennium. *Science*, 2010, **328**, 486–489.
22. Ram, S., Tree growth–climate relationships of conifer trees and reconstruction of summer Palmer Drought Severity Index at Pahalgam in Srinagar, India. *Quaternary Int.*, 2012, **254**, 152–158.

ACKNOWLEDGEMENTS. We thank Prof. Ravi S. Nanjundiah (Director, IITM, Pune) and R. Krishnan (IITM, Pune) for support and for providing infrastructure facilities. We also thank CRU for making available climatic data on the website. We also thank Forest Department staff for their cooperation during field work of trees core samples collection.

Received 28 August 2015; revised accepted 14 February 2017

doi: 10.18520/cs/v112/i12/2453-2455