

**Revised version\_Clean**

**Population differentiation of wheat leaf rust fungus *Puccinia triticina* in South Asia**

Pramod Prasad<sup>a</sup>, S.C. Bhardwaj<sup>a\*</sup>, O. P. Gangwar<sup>a</sup>, Subodh Kumar<sup>a</sup>, Hanif Khan<sup>a</sup>, Shravan Kumar<sup>b</sup>, H. C. Rawal<sup>b</sup> and T.R. Sharma<sup>b</sup>

**Affiliations:**

<sup>a</sup>ICAR-Indian Institute of Wheat and Barley Research, Flowerdale, Shimla, Himachal Pradesh, India, 171002

<sup>b</sup>ICAR-National Research Centre on Plant Biotechnology, Pusa Campus, New Delhi, India, 110012

\*Corresponding author

**S.C. Bhardwaj**

Email: [scbfdl@hotmail.com](mailto:scbfdl@hotmail.com)

Mobile: +91 94181 29900

Fax: +91 177 2620108

## **Abstract**

Leaf or brown rust caused by *Puccinia triticina* (*Pt*) is one of the most important diseases of wheat. Among the rusts, it is the most ubiquitous in all the wheat growing regions and cause considerable yield loss. Microsatellite markers based genotyping and virulence based phenotyping of 48 pathotypes of *Pt* was attempted. The pathotypes exhibit low virulence frequencies for Indian leaf rust differentials *Lr24*, *Lr9*, *Lr10*, *Lr19*, *Lr28* and *Lr9*. Using avirulence/virulence formula six major clusters of pathotypes were observed, revealing high degree of genetic variation. Molecular analysis performed using SSR markers showed high genetic diversity among the pathotypes and grouped them in seven major clusters. The percentage of polymorphic loci was ranging from 17.95 to 84.62%, Nei's gene diversity from 0.07 to 0.32 and Shannon's information index values from 0.11 to 0.47. Analysis of molecular variance revealed significantly high genetic variation within *Pt* population. Mantel's Z test proved low positive correlation ( $r=0.28$ ) between virulence and molecular diversity, suggesting independent nature of the duo. These findings offer valuable information for framing suitable disease management strategies through appropriate region specific gene deployment and improve the understanding of the population biology and evolution of *Pt* in Indian subcontinent.

**Key Words:** Leaf rust, *Puccinia triticina*, Genetic differentiation, Microsatellites, Virulence phenotype

Wheat, the most important cereal as protein source and next to rice as source of calories for majority of population in developing countries, occupy about 225 million ha area worldwide<sup>1</sup>. It fulfills 21% of the food calorie and 20% of the protein requirements of over 4.5 billion people living in more than 90 developing countries<sup>2</sup>. The predictable demand for wheat is estimated to

upsurge by 60% by 2050 in developing countries; on the other hand, wheat production is expected to reduce by 20-30% due to climate change-induced biotic and abiotic stresses<sup>3</sup>. Among the fungal diseases, rusts are the most detrimental to wheat world-wide. Brown (leaf) rust (*Puccinia triticina* Eriks.) probably causes more damage than any other rust of wheat<sup>4</sup> and is the most widely distributed disease in India<sup>5</sup>. Like other two rusts of wheat, leaf rust is preferably controlled by genetic resistance besides using fungicides and agronomic practices. However, changes in the pathogen population brought about by mutation, somatic recombination and through selection of virulent types on the major (R) genes deployed on larger areas results in the evolution of new pathotypes of the pathogen. Thus, the emergence of new pathotypes and shift in virulence patterns, render the resistant wheat varieties susceptible. For effective management of leaf rust of wheat, identification of suitable resistance sources and appropriate gene deployment strategy based on racial pattern of a particular region needs to be developed.

The degree and distribution of phenotypic and genotypic variation within and among the pathogen populations is an important aspect for understanding their population biology. Genetic structure can be used to deduce the impact of different forces that influence the evolution of a pathogen population. This in turn provides a better understanding of evolutionary pattern that may allow prediction of the potential for pathogen populations to evolve in agricultural ecosystems<sup>6</sup>. Higher genetic diversity in pathogen population within smaller area advocates the probability of rapid adaptation by the pathogen to changing host or environmental factors. On the other hand, higher genetic similarity among pathogen populations between widely separated regions could be the result of substantial long-distance dispersal of the pathogen or gene flow. Subsequently it poses a risk in deployment of disease resistance genes arrayed to local pathogen populations; as exotic pathogen, having different virulence genes may overcome resistance in

local host cultivars<sup>7</sup>. In spite of differential varieties, discrimination and changes in populations of pathogenic fungi have been detected using different molecular markers like random amplified polymorphic DNA (RAPD), restriction fragment length polymorphism (RFLP), Amplified fragment length polymorphism (AFLP) and simple sequence repeat (SSR)<sup>8,9,10</sup>. Microsatellites {simple sequence repeat (SSR)} are tandemly repeated DNA sequence units of 1–6 bp. They have abundant and random distribution throughout the genome. The SSR markers are highly polymorphic, multi-allelic, co-dominant, PCR based, reproducible and leads to high genetic diversity due to their higher mutation rate than other regions of DNA<sup>11</sup>. Recently use of microsatellites have been encouraged because of several advantages associated with these, *viz.*, the genetic markers are inherited in a Mendelian fashion as co-dominant markers. Moreover, extensive distribution throughout the organism's genome and frequent polymorphism rates make SSRs one of the most widely accepted genetic markers for population genetics studies<sup>12</sup>. Despite several benefits there are few drawbacks of using SSR markers such as their high mutation rate, the marker site may not be conserved between two species. These shortcomings could lead to misleading interpretation for population structure studies. To overcome such gaps some new markers such as the single-nucleotide polymorphism (SNP) markers; which can detect variation in a single nucleotide that occurs at a specific position in the genome, are promising for future research on population structure. On the presumption that knowledge of the population biology of *Pt* could eventually contribute to the development of more durable disease management strategies, the present study was conducted with the hypothesis that molecular polymorphism in South Asian collection of *Pt* population is dependent on their differential virulence and geographical distribution. Genotypic variation among leaf rust pathotypes from South Asia was explored using microsatellites.

## **Materials and Methods**

### *Pathogen isolation, purification and multiplication*

For genetic differentiation studies 2980 *Puccinia triticina* isolates were collected from different parts of South Asia. Indian wheat leaf rust differential set<sup>13</sup> (Table 1) categorized these isolates in forty eight pathotypes (Table 2) which are being maintained at National repository of wheat rust pathotypes at Regional Station of ICAR-Indian Institute of Wheat and Barley Research (erstwhile Directorate of Wheat Research), Shimla. Isolation, purification and multiplication of pathotypes were done on susceptible wheat cv. Agra Local; except for pathotypes 16 and 16-1, which were multiplied on Khapli, a Diccocum wheat. The sterilized mixture of fine loam and farmyard manure (3:1) was used for sowing of plant material. Five grams of nitrogen (N):phosphorus (P):potassium (K) (12:32:16) fertilizer mixture per 5kg of soil was used as nutrient supplement. The seedlings were raised in plastic pots (12.5cm diameter) inside the spore-proof chambers (indoors) at  $22\pm 2^{\circ}\text{C}$  and 50-70% relative humidity along with 12-hour daylight. When seedlings were ten days old, 15 ml of maleic hydrazide acid (2.5%) was added to soil in each pot before inoculation to reduce the aging of seedling and encourage sporulation. One week old seedlings were atomized with uredosporic inoculum suspension in non-phytotoxic isoparaffinic mineral oil Soltrol 170 (Chevron Phillips Chemical Company, US) and incubated in saturated humidity chambers for 48 hours. Subsequently these plants were transferred on to the greenhouses where temperature of  $20\pm 2^{\circ}\text{C}$  and relative humidity of 50-70% was maintained. Inoculum collected after fifteen days of inoculation was used to test the purity of the pathotypes. Purity of the pathotypes and avirulence/virulence structure were confirmed after examining the characteristic infection type of the isolates (pathotype) on each of the entries of differential set. After ensuring the purity of all the pathotypes they were mass multiplied on susceptible host as mentioned earlier. During the whole multiplication procedure, pots inoculated

with different pathotypes were kept in separate cabins for ensuring pathotype purity. After 15 days of inoculation the uredospores were collected, dried in a desiccator at 4<sup>0</sup>C for three days and stored at -20<sup>0</sup>C till further use.

#### *Virulence diversity*

Virulence diversity of the pathotypes was determined using the Indian leaf rust differential (Table 1) sets for pathotype nomenclature, consisting of 9, 9 and 7 entries in subsets ‘0’, ‘A’ and ‘B’, respectively<sup>13,14</sup>. Uredospores from all the pathotypes were inoculated on ten days old seedlings of differential set. After inoculation, seedlings were placed in incubation chambers for 48 h at >80% relative humidity and then transferred into a greenhouse and kept at 20±2<sup>0</sup>C for 15 days. Based on the presence of necrosis and chlorosis and the intensity of sporulation, infection type of each pathotype was scored individually on each entry. Infection types (IT) score based on a 0-4 rating scale was followed<sup>15</sup>. On standard evaluation scale infection types 0-2 (immune to moderate uredia with necrosis and/or chlorosis) was considered low infection types or avirulent and 3-4 (moderate to large uredia without chlorosis or necrosis) were considered high infection types or virulent.

#### *Molecular analysis of P. triticina*

##### *DNA extraction*

Forty eight *Pt* pathotypes, used in the virulence analysis, were studied for molecular genotyping. Uredospores were harvested from infected leaves 15 days post inoculation, dried and kept at -20<sup>0</sup>C. Total genomic DNA from dried uredospores (~50 mg) was extracted with cetyltrimethylammonium bromide (CTAB) as described by Kiran *et al.*<sup>16</sup>. The purity of genomic DNA and quantification of template DNA for PCR was measured in duplicate using NanoDrop 2000® UV-Vis Spectrophotometer (Thermo Scientific) and stored at -20<sup>0</sup>C till further use.

### *SSR analysis*

*Pt* specific simple sequence repeat (SSR) primers (Table 3) were designed from the genome sequences of *P. triticina* sequenced in our previous study<sup>16</sup> using Primer3 software and synthesized from Agile Life Science Technologies India Pvt. Ltd. All the PCR reactions were carried out in 20  $\mu$ L volume containing 25 ng of template DNA, 200  $\mu$ M each of the four dNTPs, 1X PCR buffer (10 mM Tris pH 9.0, 50 mM KCl), 1.5 mM MgCl<sub>2</sub>, 0.5 U *Taq* polymerase (Himedia Laboratory Pvt. Ltd., India) and 10 pmol of both forward and reverse primer. The reaction programs were set at 94<sup>0</sup>C for 2 min, followed by 35 cycles of 30 sec at 94<sup>0</sup>C, 30 sec at a primer annealing temperature and 1 min at 72<sup>0</sup>C, with a final extension at 72<sup>0</sup>C for 10 min in thermal cycler (Boeco thermal cycler TC-PRO). After completion of the amplification, the amplified DNA was analyzed on 3% Super MT4 agarose gel (Life Technologies, India, Pvt. Ltd.) in 1X TBE buffer at 65–70 V for 2-3 h. DNA fragments were visualized under the UV light and photographed using Gel Documentation System (Bio-Rad Laboratories, Inc.).

### *Data analysis*

Virulence frequency was determined as percentage of the pathotypes virulent for specific gene or entry of the differential set from total of pathotypes under study. The virulence and avirulence infection types (ITs) of each isolate on the differential genotypes were assigned with binary codes of 1 and 0, respectively. The presence or absence of individual, distinct, and reproducible bands was scored as '1' for presence and '0' for absence. Binary data were used to calculate Jaccard similarity coefficient. Cluster analysis was performed using NTSYSpc version 2.0<sup>17</sup> and dendrogram was constructed using the unweighted pair group method with arithmetic average (UPGMA).

Genetic diversity parameters including the observed number of alleles (Na), the effective number of alleles (Ne), the percentage of polymorphic loci (Pp), Nei's gene diversity (h),

Shannon's index (I) were calculated to estimate the level of genetic variation using POPGENE version 1.31<sup>18</sup>. Analysis of molecular variance (AMOVA) was performed using ARLEQUIN version 3.0<sup>19</sup> to examine differences among and within *Pt* populations. STRUCTURE 2.3.4 software<sup>20</sup> was used to analyse population structure using a burn-in period of 10,000 and 100,000 Markov Chain Monte Carlo (MCMC) replications. Mantel's Z test in zt version 1.1<sup>21</sup>, was used to test correlation between molecular genotypes and virulence phenotypes.

## Results

### *Virulence diversity and cluster analysis*

Avirulence/virulence patterns based on data from the Indian leaf rust differentials were determined separately for all the pathotypes (Table 4). Virulence frequency of forty eight South Asian *Pt* pathotypes varied from 0.00 (for *Lr24*) to 95.8 % (for Agra Local) (Figure 1). The pathotypes displayed low virulence frequencies (less than 4 %) for Kharchia Mutant (*Lr9+*), Raj1555, *Lr19*, *Lr28* and HP1633. The virulence frequency of pathotypes was moderate (20-43%) to IWP 94, Raj 3765, PBW 343, UP 2338, K 8804, HD 2189, *Lr15*, *Lr10*, and Benno (*Lr26*) and high (45-95%) to Agra Local, *Lr14a*, *Lr18*, *Lr13*, *Lr17a*, Loros (*Lr2c*), Webster, Democrat, Thew (*Lr20*) and Malakoff (*Lr1*). None of the pathotypes was virulent to *Lr24*.

Virulence data generated six major clusters of the pathotypes showing 48 to 66% similarity within the cluster (Figure 2). Clusters B, C and D were further divided in to sub clusters. Cluster E and F had only one pathotype each i.e. 16-1 and 16, respectively. These two pathotypes have unique character of their virulence on Raj1555 and avirulence on Agra Local which separate them from rest of the pathotypes. Pathotype 16-1, showing virulence to leaf rust resistance genes *Lr14a*, *Lr18*, *Lr2c* (Loros) and *Lr20* (Thew) differ from pathotype 16, which is avirulent to these genes. Pathotypes 77-1, 77-2, 77-3, 77-4, 77-8, 77A and 77A-1, virulent on *Lr14a*, *Lr13*, *Lr17a*, *Lr15*, *Lr10*, *Lr2c*, Webster, Democrat and Malakoff (*Lr1*) and avirulent on



*Lr24*, *Lr28*, and HP 1633, shared more than 75% virulence similarity. Pathotypes 11 and 63 displayed 100% similarity.

#### *Population genetic variation*

The percentage of polymorphic loci recorded in the *Pt* populations from twelve states was 97% (Table 5). Isolates collected from Karnataka had the highest percentage of polymorphic loci (84%) followed by Tamil Nadu (82%), while the isolates from Uttarakhand had the lowest percentage (17%) followed by Punjab (20%). The observed alleles per locus were highest for Karnataka (1.85) and lowest for Uttarakhand (1.18). However, the effective number of alleles was highest for pathotypes collected from Maharashtra (1.8) followed by Tamil Nadu (1.56). The average effective number of alleles was 1.59 with Uttarakhand (1.12), Punjab (1.15) and Uttar Pradesh (1.18) having lesser effective number of alleles. Nei's genetic diversity for whole *Pt* population was 0.34, with Tamil Nadu (0.32) and Uttarakhand (0.07), having the highest and lowest values, respectively. Highest Shannon's Information index (SII) was observed among the pathotypes from Tamil Nadu (0.47), whereas the lowest SII was for the pathotypes from Uttarakhand (0.11). The overall total variability, variability within population, diversity among populations and Fixation Index was found to be 0.33, 0.17, 0.52 and 0.13, respectively. Analysis of AMOVA ( $P < 0.001$ ) convened 87.43% genetic variation within the *Pt* population, whereas the variation among populations was 12.57% (Table 6).

#### *Molecular genotyping*

Among the primers (Table 3) used, SSR-PTCCG-36, SSR-PTGGA-32, SSR-PTATTG-60, SSR-PTCTTT-50, SSR-PCCCGT-35, SSR-PGTGGA-35, SSR-PTGAGGA-48, SSR-PCCAGAA-48 and SSR-PGCTGTT-60 were found polymorphic. These nine primers amplified 45 alleles with an average of 5 alleles per primer. Primer SSR-P TCTTT-50 and SR-PTGGA-32 amplified highest (11) and lowest (2) number of alleles, respectively. Interestingly, the allelic

pattern with primer SSR-PTATTG-60 was more or less similar among the pathotypes of virulence based groups (Supplementary Figure). Primer SSR-PTCTTT-50 amplified maximum alleles followed by SSR-PTATTG-60 and SSR-PTGAGGA-48 (Table 3).

The overall molecular marker data revealed poor genotypic similarity among pathotypes. The maximum genotypic similarity (89%) was observed between pathotypes 12-3 and 12-7. This was followed by 87% genotypic similarity among pathotypes 12-3, 12-6 and 12-7. Cluster analysis generated seven major clusters (Figure 3). Cluster A and E had single pathotype 16 and 104-2, respectively. The results of STRUCTURE 2.3.4 analyses performed on *Pt* population data set indicated two distinct subpopulations (K) (Figure 4). The subpopulations S1, S2 and M contained fifteen, twenty six and seven pathotypes each of *Pt*, respectively (Supplementary file). The genetic distance as fixation index (FST) within S1 and S2 subpopulations was 0.3275 and 0.3086, respectively.

#### *Mantel's Z test*

Correlation analyses using the Mantel's *Z* test revealed positive but weak association between molecular and virulence data. There was positive ( $r=0.28$ ) and statistically significant ( $p = 0.0007$ ) correlation between molecular and virulence data. The phylogenetic trees did not indicate such relationship between SSR genotypes and virulence phenotypes.

### **Discussion**

This study was undertaken to distinguish the phenotypic and genotypic variations in *Pt* populations collected from different parts of South Asia. The result suggests that populations of *Pt* are highly variable and majority of genetic variation was distributed within population (Table 6). This is the first report on the diversity and population dynamics of *Pt* infecting wheat in South Asia. Virulence study revealed that none of the pathotype was virulent on *Lr24*. Although

several wheat varieties carrying effective leaf rust resistance gene *Lr24* (linked to *Sr24*) occupy relatively larger proportion of cultivated varieties in South Asia, yet wheat breeders need to diversify leaf rust resistance as presence of *Lr24* in many varieties poses a potential threat of boom and bust cycle due to emergence of *Lr24* virulence in South Asia. Moreover, there is possibility of entry of *Lr24* virulence from neighbouring countries<sup>22</sup>. Virulence frequency for *Lr14a*, *Lr18*, *Lr13*, *Lr17a*, *Loros (Lr2c)*, *Webster*, *Democrat*, *Thew (Lr20)* and *Malakoff (Lr1)* was on the higher side. Therefore, discouraging cultivation of cultivars carrying these genes would definitely affect the survival of the pathotypes which are virulent on these genes/entries. The virulence based clustering suggests that the pathotypes were highly variable in their virulence to Indian leaf rust differentials. Up to some extent, the pathotypes belonging to similar virulence group (group 77 and 12) were also grouped in virulence based phylogenetic tree (Figure 2). Pathotypes 16 and 16-1 form two separate major groups in the virulence based phylogenetic tree, which is obvious from their different avirulence/virulence nature from rest of the pathotypes. In spite of lacking a sexual recombination cycle in Indian sub-continent<sup>23</sup> the high genetic diversity of the leaf rust pathogen up to certain extent may be explained by the phenomenon like introduction of exotic and genetically distinct pathotypes, recurrent mutation etc., that are reported to be responsible for variability in wheat rust pathotypes<sup>24</sup>.

Molecular marker data based clustering revealed high variability among the pathotypes. A total of seven major clusters were observed. Molecular markers based polymorphism was relatively higher as compared to virulence based polymorphism. The genotypic clustering pattern of pathotypes was quite random as compared to virulence based clustering as there was no uniform grouping among the pathotypes belonging to one particular geographical region or virulence based groups. This could be justified through the highly migratory nature of *Pt* and

virulence independent nature of microsatellite markers used in the study. Clustering based on virulence and molecular analysis was poorly correlated to each other. This is quite understandable from the fact that pathogenicity or virulence in the pathogen is not at all related to the molecular markers like RAPD, SSRs and other markers unless these markers are designed from the part of the genome of the pathogen which decides the pathogenicity or virulence<sup>25,26</sup>. Primers SSR-PTATTG-60 and SSR-PTCTTT-50 supported the virulence based clustering as they displayed almost similar allelic patterns among the pathotypes of virulence based groups. This statement can be supported by assuming that these primers may have amplified the parts of the genome having sequences responsible for pathogen virulence<sup>26</sup> but we cannot assure it unless further functional studies of these markers are undertaken. Pathotypes 12-1, 12-3, 12-6, 12-7 and 12-8 shared more than 80% similarity, likewise pathotypes 77-2, 77-3, 77-6, 77A and 77A-1 shared more than 70% similarity, which indicated that these pathotypes might have evolved from genetically similar ancestors<sup>27</sup>. The effective number of alleles, Nei's gene diversity and Shannon's Information index was highest for pathotypes from Tamil Nadu. Green bridges or cultivation of wheat crop round the year may be helping the mutated isolates in their survival in higher hills of Tamil Nadu and thereby diversifying *Pt* population. Moreover; it is advocated that Nilgiri and Palni hills in Tamil Nadu, which are the primary foci as the source of brown rust pathotypes for Tamil Nadu and Karnataka<sup>23</sup>, receives high UV intensity sunlight. The higher UV intensity in these hills might be contributing towards faster evolution of rust pathogens through mutation. Very short duration of congenial environment for wheat leaf rust development in north Indian states followed by harsh summers do not allow the mutated isolates of *Pt* to adopt such conditions and survive until the next crop season, which could be a reason for lesser Nei's gene diversity in this study for *Pt* population from Uttarakhand (0.07) and Punjab (0.08).

Analysis of molecular variance (AMOVA) revealed 87.43% variation within the *Pt* populations from different states. Such variation within the *Pt* populations indicates that they have the potential to evolve relatively quickly to changing climate and resistance pattern. However; the genetic variation of *Pt* populations among the populations was mere 12.57%, suggesting similarity between molecular genotypes from distant areas. Uredospores of *Pt* as other rust pathogens are wind-dispersed, and circumstantial evidence of migration over hundreds or thousands of miles has been reported<sup>28</sup>. This migratory nature of the pathogen may be responsible for the limited genetic variation between two distant populations observed in this study. Similar findings are also reported by Hovmoller *et al.*<sup>29</sup>, where they observed similarity between virulence patterns and AFLP data among the *Pt* populations from United Kingdom, France, Germany, and Denmark. STRUCTURE programme differentiated *Pt* population into three subpopulations S1, S2 and M. More than 50% of the pathotypes were grouped in subpopulation S2. The *Fst* values for subpopulations S1 and S2 was more than 0.25, which belongs to very great genetic differentiation category<sup>30</sup> and corroborates with high genotypic and virulence variations, and diverse geographic distribution of the *Pt* pathogen subpopulations. Using STRUCTURE version 2, similar significant variation in *Pt* population from Central Asia and the Caucasus was also observed, which separated these population from the North American isolates of *Pt*<sup>31</sup>.

Mantel's *Z* test suggests that molecular diversity is poorly co-related to the virulence diversity among *Pt* pathotypes. This was quite obvious as such correlation would have appeared if we had used virulence specific primers but here microsatellites were used, which shows virulence or pathogenicity independent DNA polymorphism. Chen *et al.*<sup>32</sup> justified this fact by reporting that molecular polymorphism was independent of pathogenicity and whole genome of

the pathogen evolves at a much faster rate than genes governing pathogenicity in yellow rust of wheat (*P. striiformis*).

In conclusion, South Asian collection of *Pt* was highly variable for virulence phenotype and SSR genotypes. They were placed in six virulence phenotype and seven SSR genotypes based groups. This high diversity among the pathotypes might be the result of evolution and mutation in the pathogens along with the long-term cultivation of different wheat varieties in different wheat growing zones of the country. Moreover, the introduction of new virulence from neighbouring countries may be the other factor contributing to the high pathotype diversity. The virulence and molecular based variability studies of wheat leaf rust pathotypes from different South Asian countries might provide some versatile information on origin and further movement of new virulence among the countries. The finding of this study would provide a reference for wheat leaf rust resistance breeding, a better understanding of *Pt* population dynamics, a preliminary idea for designing breeding strategies at regional level, a scientific awareness of deploying available resistance sources for disease management and assist in tracking variation in *Pt* population over time and space.

#### ACKNOWLEDGEMENTS

We thank Department of Biotechnology, Govt. of India for financial support. We also thank Indian Council of Agricultural Research, New Delhi and Director, ICAR-Indian Institute of Wheat and Barley Research (ICAR-IIWBR), Karnal for providing institutional infrastructure and facilities to conduct experiments.

#### **Funding**

This work was supported by the Department of Biotechnology, Govt. of India via grant number BT/PR13631/AGR/02/717-C/2010 dated 20.09.2010.

**Conflict of Interest:** The authors declare that they have no conflict of interest.

## References

1. Bhardwaj, S. C., Prashar, M. and Prasad, P., Ug99-Future Challenges. In Future Challenges in Crop Protection (eds. Goyal, A. and Manoharachary, C.), Springer Science and Business Media, New York, USA, 2014, pp. 231-247.
2. Braun, H. J., Atlin, G. and Payne, T., Multi-location testing as a tool to identify plant response to global climate change. In Climate Change and Crop Production (ed. Reynolds, M. P.), CABI, London, UK, 2010, pp. 115–38.
3. Rosegrant, M. W. and Agcaoili, M., Global food demand, supply, and price prospects to 2010. International Food Policy Research Institute, Washington, D.C. USA, 2010.
4. Herrera-foessel, S. A. *et al.*, New slow-rusting leaf rust resistance genes *Lr67* and *Yr46* in wheat are pleiotropic or closely linked. *Theor. Appl. Genet.*, 2011, **122**, 239-249.
5. Bhardwaj, S. C., Prashar, M., Jain, S. K., Kumar, S. and Sharma, Y. P., Physiologic specialization of *Puccinia triticina* on wheat (*Triticum* species) in India. *Indian J. Agric. Sci.*, 2010, **80**(9), 805-11.
6. McDonald, B. A., The population genetics of fungi: tools and techniques. *Phytopathology*, 1997, **87**, 448–53.
7. Brasier, C. M., Rapid changes in genetic structure of epidemic populations of *Ophiostomaulmi*. *Nature*, 1988, **332**, 538–541.
8. Kashyap, P. L., Rai, S., Kumar, S. and Srivastava, A. K., Genetic diversity, mating types and phylogenetic analysis of Indian races of *Fusarium oxysporum* f. sp. *ciceris* from chickpea. *Arch. Phytopathology Plant Protect.*, 2016, **49**, 533-553.
9. Kolmer, J. A., Tracking wheat rust on a continental scale. *Curr. Opin. Plant. Biol.*, 2005, **8**, 1-9.
10. Ordonez, M. E. and Kolmer, J. A., Simple sequence repeat diversity of a world-wide collection of *Puccinia triticina* from durum wheat. *Phytopathology*, 2007, **97**, 574–583.
11. Singh, R., Kumar, S., Kashyap, P. L., Srivastava, A. K., Mishra, S., Sharma, A. K., Identification and characterization of microsatellite from *Alternaria brassicicola* to assess cross-species transferability and utility as a diagnostic marker. *Mol. Biotechnol.*, 2014, **56**(11), 1049-1059.
12. Morgante, M., Hanafey, H. and Powell, W., Microsatellites are preferentially associated with non repetitive DNA in plant genome. *Nature Genet.*, 2002, **30**(2), 194–200.

13. Nagarajan, S., Nayar, S. K. and Bahadur, P., The proposed brown rust of wheat (*Puccinia recondita* f.sp. *tritici*) virulence monitoring system. *Curr. Sci.*, 1983, **52**(9), 413-416.
14. Bhardwaj, S. C., Gangwar, O. P., Singh, S. B., Saharan, M. S. and Sharma, S., Rust situation and pathotypes of *Puccinia* species in Leh Laddakh in relation to recurrence of wheat rusts in India. *Indian Phytopathol.*, 2012, **65**(3), 230-232.
15. Stakman, E. C., Stewart, D. M., Loegering, W. Q., Identification of physiologic races of *Puccinia graminis* var. *tritici*. U.S, Agricultural Research Service, ARS E617, 1962, p. 1-53.
16. Kiran, K., Rawal, H. C., Dubey, H., Jaswal, R., Devanna, B. N., Gupta, D. K., Bhardwaj, S. C., Prasad, P., *et al.*, Draft genome of the wheat rust pathogen (*Puccinia triticina*) unravels genome-wide structural variations during evolution. *Genome Biol. Evol.*, 2016, **8**(9), 2702–2721. doi:10.1093/gbe/evw197.
17. Rohlf, F. J., NTSYS-pc: numerical taxonomy and multivariate analysis system, version 2.1. Exeter Software: Setauket, NY, 2000.
18. Yeh, F. C., Yang, R. C. and Boyle, T., POPGENE VERSION 1.31: Microsoft Window-based free Software for Population Genetic Analysis, ftp://ftp.microsoft.com/Softlib/HPGL.EXE, 1999.
19. Excoffier, L., Laval, G. and Schneider, S., Arlequin (version 3.0): an integrated software package for population genetics data analysis. *Evol. Bioinform.*, 2007, **23**(1), 47-50.
20. Pritchard, J. K., Stephens, M. and Donnelly, P., Inference of population structure using multilocus genotype data. *Genetics*, 2000, **155**, 945–959.
21. Bonnet, E. and Van de Peer, Y., zt: a software tool for simple and partial Mantel tests. *J. Stat. Softw.*, 2002, **7**, 10.
22. Mishra, A. N., Prakasha, T. L., Kaushal, K. and Dubey, V. G., Validation of *Lr24* in some released bread wheat varieties and its implications in leaf rust resistance breeding and deployment in central India. *Indian Phytopathol.*, 2014, **67**(1), 102-103.
23. Mehta, K. C., Further studies on cereal rusts of India, Part II. Scientific Monograph.18, ICAR, New Delhi, 1952.
24. Hovmoller, M. S., Yahyaoui, A. H., Milus, E. A. and Justesen, A. F., Rapid global spread of two aggressive strains of a wheat rust fungus. *Mol. Ecol.*, 2008, **17**, 3818–3826.
25. Kolmer, J. A. and Acevedo, M. A., Genetically divergent types of the wheat leaf fungus *Puccinia triticina* in Ethiopia, a center of tetraploid wheat diversity. *Phytopathology*, 2016, **106**(4), 380-385.



26. Koch, E., Song, K., Osborn, T. C. and Wellings, P. H., Relationship between pathogenicity and phylogeny based on restriction fragment length polymorphism in *Leptosphaeria maculans*. *MPMI*, 1991, **4**, 341–349.
27. Bhardwaj, S. C., Wheat rust pathotypes in Indian subcontinent then and now. In Wheat-Productivity enhancement under changing climate (eds. Singh, S. S., Hanchinal, R. R., Singh, G., Sharma, R. K., Saharan, M. S. and Sharma, I.), Narosa Publishing House, New Delhi, India, 2012, pp. 227-238.
28. Nagarajan, S., Singh, H., Joshi, L. M. and Saari, E. E., Meteorological conditions associated with long distance dissemination and deposition of *Puccinia graminis tritici* uredospores in India. *Phytopathology*, 1976, **66**, 198-203.
29. Hovmoller, M. S., Justesen, A. F. and Brown, J. K. M., Clonality and long-distance migration of *Puccinia striiformis* f. sp. *tritici* in northwest Europe. *Plant Pathol.*, 2002, **51**, 24-32.
30. Hartl, D.L. and Clark, A.G., Principles of Population Genetics, 3rd edn. Sinauer Associates, Inc, Sunderland, MA, 1997.
31. Kolmer, J. A. and Ordonez, M. E., Genetic differentiation of *Puccinia triticina* populations in Central Asia and the Caucasus. *Phytopathology*, 2007, **97**, 1141-1149.
32. Chen, X. M., Penman, L., Wan, A. M. and Cheng, P., Virulence races of *Puccinia striiformis* f. sp. *tritici* in 2006 and 2007 and development of wheat stripe rust and distributions, dynamics, and evolutionary relationships of races from 2000 to 2007 in the United States. *Can. J. Plant Pathol.*, 2010, **32**(3), 315-333.

**Table 1** Leaf rust pathotypes of wheat (*Puccinia triticina*) studied for virulence and molecular diversity

Name of the pathotype <sup>Y</sup>	Year of first detection/collection	Place of collection	First detected on (Host)	Susceptible <i>Lr</i> genes/ Lines
10	1931	Punjab	Local	Malakoff
11	1947	Multai, Madhya Pradesh	A090	20
12	1966	Thordi, Gujarat	Local	3
12-1	1983	Gwalior, Madhya Pradesh	Local	26
12-2	1979	Hansi, Haryana	Local	23
12-3	1989	Dharwad, Karnataka	Bijaga Yellow	15, 26
12-4	1990	Pusa, Bihar	Agra Local	3, 10, 20
12-5	2005	Hubli, Karnataka	Local Red	23, 26
12-6	2006	Dharwad, Karnataka	KH65	20, 26
12-7	2008	Arabhavi, Karnataka	HP1912	10, 23, 26
12-8	2008	Pantnagar, Uttarakhand	HD2204	15, 20, 26
16	1959*	Nilgiris, Tamil Nadu	Khapli	Agra local**
16-1	2004	Belgaum, Karnataka	Dicoccum	Agra local, Raj1555
17	1957	Nilgiris, Tamil Nadu	Mal-4	Malakoff, 15, 20
20	1935	Hisar, Haryana	UP Local	Malakoff, 20
63	1931	Shimla, Himachal Pradesh	Local	20
77	1954	Pusa, Bihar	Mal 4	3, 15
77-1	1985	Nilgiris, Tamil Nadu	Burgas-2	20, 26
77-2	1984	Nilgiris, Tamil Nadu	PAU Wheat	20, 23
77-3	1989	Nilgiris, Tamil Nadu	<i>Lr15</i>	26
77-4	1989	Nilgiris, Tamil Nadu	Lal Bahadur	23
77-5	1992	Nilgiris, Tamil Nadu	Crossing material	23, 26
77-6	1997	Nilgiris, Tamil Nadu	C306	23, 26
77-7	1998	Nilgiris, Tamil Nadu	<i>Lr9</i>	9, 23, 26
77-8	2004	Arabhavi, Karnataka	Off type in Kh. mutant	19
77-9	2008	Belgaum, Karnataka	HI1511	23, 26/ 2**
77-10	2008	Nilgiris, Tamil Nadu	HW5212	23, 26,28
77-11	2009	Ugar, Karnataka	HD2932	23, 20/ 26**
77-12	2013	Wellington, Tamil Nadu	Local	23,26/2a,2c,20**
77 A	1974	Dharwad, Karnataka	CC62	10, 20
77A-1	1976	Haryana	HD 2009	20**
104	1973	Janakpur, Nepal	Local	1, 3
104-1	1985	Borkheda, Maharashtra	Kalyansona	20,23
104-2	1991	Malan, Himachal Pradesh	Off type in Transec	23, 26 / 20**

104-3	1993	Naval ParasiSunwal, Nepal	local wheat & <i>Lr20</i>	20, 23, 26
104-4	2010	Arki, Himachal Pradesh	HS240	HS240/ Democrat**
104A	1975	Patna, Bihar	HD1981	20
104B	1980	Hansi, Haryana	C306	23
106	1935	Haldwani, Uttarakhand	Local	2c, 20
107	1935	Khanewal, Punjab	PB8A	15
107-1	1988	Dharwad, Karnataka	Local	15, 26
108	1935	Banaras, Uttar Pradesh	Pusa 4	Malakoff, 20
108-1	1989	Deorighat, Himachal Pradesh	Sonalika	Malakoff, 3, 15, 20
162	1957	Gwalior, Madhya Pradesh	NP720	10
162-1	2002	Akola, Maharashtra	Agra Local	10, 20, 26
162-2	2005	Kanpur, Uttar Pradesh	Agra Local	10, 26
162-3	2006	Ahmadnagar, Gujarat	GW1	10**
162A	1961	Nilgiris, Tamil Nadu	Local	10, 20
<sup>†</sup> based on old system (well-known globally) of Indian wheat rust nomenclature, names based on new system are presented in supplementary file. * Replaced in 2000, ** Avirulent				

**Table 2** Set of differentials for the identification of pathotypes of *Puccinia triticina* on wheat in India

<b>Set 0</b>	<b>Set A</b>	<b>Set B</b>
IWP 94 ( <i>Lr23+</i> )	<i>Lr14a</i>	Loros ( <i>Lr2c</i> )
Kharchia Mutant ( <i>Lr9</i> )	<i>Lr24</i>	Webster ( <i>Lr2a</i> )
Raj 3765 ( <i>Lr13+10+</i> )	<i>Lr18</i>	Democrat ( <i>Lr3</i> )
PBW 343 ( <i>Lr26</i> )	<i>Lr13</i>	Thew ( <i>Lr20</i> )
UP 2338 ( <i>Lr26+34+</i> )	<i>Lr17</i>	Malakoff ( <i>Lr1</i> )
K 8804 ( <i>Lr26+23+</i> )	<i>Lr15</i>	Benno ( <i>Lr26</i> )
Raj 1555	<i>Lr10</i>	HP1633 ( <i>Lr9+</i> )
HD 2189 ( <i>Lr13+34+</i> )	<i>Lr19</i>	
Agra Local	<i>Lr28</i>	

**Table 3** Markers used in genetic diversity study of leaf rust isolates

Primer code	Forward primer sequence (5' - 3')	Reverse primer sequence (5' - 3')	AT* (°C)	Number of amplified alleles	Allele size (kb)
SSR-P GT-42	GGGGTGAGTTTCTGTATTGA	CAGAGATCATCGAGGAAAAC	50.4	02	0.08-0.3
SSR-P AG-40	CTTTCTTACCCCCACAACACTAC	CTCTCTCTCTCTCTCTCTCTCTC	53	01	0.12
SSR-P CT-36	ACTCTCAAACACTCACTCCCTCT	GACTACACCATTTCAAAACCAA	51.9	01	0.13
SSR-P AC-32	ACAAAACAAAACAGATCCACTG	ACGTATTTGGTCTTCTTCTCC	51	01	0.13
SSR-P TC-32	TAGAATTCTTGGTAGGACGAG	CGGTCAGAGTGTCTGTCAATA	50	02	0.12-0.3
SSR-P CAA-60	AACTGCGAGGACAACCTTC	CGTCTGCTGAGTTTCTGTATT	50	01	0.13
SSR-P AGA-48	CAAACGAAGCAAACCTAGAAGA	TGTTGTTGTTGTTGTTGTTGT	50	01	0.12
SSR-P GGT-45	GCTGCTTGATGGAGGATG	AACAGCTTCAGCGACCTC	51	02	0.13-0.3
SSR-P GTT-45	GATGAGGTTGTTGAAGGAGA	ACCAGAACCAACAAAACAAC	49.6	02	0.14-0.4
SSR-P CAC-45	GAAGACCATCCTCACGACT	TTCTTCTTGTGGTTTTCTG	51	02	0.13-0.3
SSR-P CAAC- 44	AGCGTAGAGTCAGTCAGTCAG	GCTAATAAGGAGATTGGGTTG	51	02	0.1-0.25
SSR-P TATC-40	AAGCGTGATCAAGTAGGTTTA	GATGGACAAGTAGAGAGATGG	50.4	01	0.1-0.3
SSR-P TCCG-36	TTTTTCTAGATCCACCAACC	TACGAACAGGAGTCCCTCA	50.4	03	0.15-1.0
SSR-P AGCC-32	GGGAAAGAAAAACACATCCT	GTCTCTTCGCTGATCTGG	50	01	0.15
SSR-P TGGA-32	GCATTTGTTTTGTTGATTG	AGACACCTCCCCTTAAAAAC	48	02	0.12-0.2
SSR-P TATTG-60	TCAAACAACCTCATCCTGAAC	ATGTGATATCTTTGGATTGG	48	10	0.15-0.5
SSR-P TCTTT-50	GGGTTTATATGGTGGGTGT	GTTGAGTGGGTGAGATGAGTA	48	11	0.15-0.5
SSR-P TAGCG-40	GCTAACGCTATGCAAAATAGA	CAGTTCAGTACCCACCAGTTA	48	02	0.12-0.35
SSR-P CCCGT-35	TTTTTGAAGGGCTTGTAGTG	AAAGGGACAGTTATGGGATAG	48	03	0.09-0.7
SSR-P GTGGA-35	TGTTTGGGAGTGTATGTGTG	GCCGAGTACCACTACCACTA	20	04	0.09-1.0
SSR-P TGAGGA-48	GTATCGGATGTTGTTGTGAAG	CTACCAAGTCTATCCGTCCTC	57.9	05	0.08-1.0
SSR-P ACAAAC-48	ATACATTTGGTTACCCACCT	TGTGTTTGTGTTGTTGTTGT	48.9	01	0.1-1.3
SSR-P CCAGAA-48F	GAAGAACTCGATCCCAGAA	CTGGTTTGTGTTGTTGTTG	49.6	04	0.15-0.9
SSR-P CCGCAC-60	TTTTGGCTGAAGTTCTGAAT	GTTGTTGAGTTGAAGGACAAG	50.9	02	0.14-0.35
SSR-P GCTGTT-60 F	GATGAGCAGCATGAGGAG	CACCAGAACAACATACTCCAT	51.9	03	0.15-0.25

\*Annealing Temperature

**Table 4** Avirulence/virulence pattern of leaf rust pathotypes on *Lr* genes

Name of Pathotype	Avirulence/Virulence pattern
10	<i>Lr</i> 3, 9, 10, 15, 17a, 17b, 19, 21, 23,24, 25, 26, 27, 28, 29, 32, 36, 39, 42, 43, 45, 47 / <i>Lr</i> 1, 2 <sup>a</sup> , 2b, 2c, 11, 12, 13, 14a, 16, 18, 22a, 22b, 30, 33, 34, 35, 37, 38, 40, 44, 48, 49
11	<i>Lr</i> 1, 2a, 2b, 2c, 3, 9, 10, 12, 13, 14a, 14b, 14ab, 15, 16, 17a, 17b, 18, 19, 21, 22a, 22b, 23, 24, 25, 26, 28, 29, 30, 32, 33, 34, 36,37, 38, 39, 40, 42, 43, 44, 45, 47, 48, 49/ <i>Lr</i> 11, 20, 27+31, 35
12	<i>Lr</i> 1, 2a, 9, 14b, 15, 17a, 17b, 18, 19, 20, 23, 24, 25, 26, 28, 29, 32, 36, 38, 39, 42, 43, 45, 47/ <i>Lr</i> 2b, 2c, 3, 10, 11, 12, 13, 14a, 14ab, 16, 21, 22a, 22b, 27, 30, 33, 34, 35, 37, 40, 44, 48, 49
12-1	<i>Lr</i> 1, 2a, 9, 10, 11, 15, 17b, 19, 20, 23, 24, 25, 28, 29, 32, 36, 39, 42, 43, 45, 47/ <i>Lr</i> 2b, 2c, 3, 12, 13, 14a, 14b, 14ab, 16, 17a, 18, 21, 22a, 22b, 26, 27, 30, 33, 34, 35, 37, 38, 40, 44, 48, 49
12-2	<i>Lr</i> 1, 2a, 9, 10, 13, 15, 17a, 17b, 18, 19, 20, 24, 25, 26, 28, 29, 32, 36, 39, 40, 42, 43, 45, 47/ <i>Lr</i> 2b, 2c, 3, 11, 12, 14a, 14b, 14ab, 16, 21, 22a, 22b, 23, 27+31, 30, 33, 34, 35, 37, 38, 44, 48, 49
12-3	<i>Lr</i> 1, 2a, 9, 19, 20, 23, 24, 25, 28, 29, 32, 36, 39, 42, 43, 45, 47/ <i>Lr</i> 2b, 2c, 3, 10, 11, 12,13,14a, 14b, 14ab, 15, 16, 17a, 17b, 18, 21, 22a, 22b, 26, 27, 30, 33, 34, 35, 37, 38, 40, 44, 48,49
12-4	<i>Lr</i> 1, 2a, 9, 15, 19, 23, 24, 25, 26, 28, 29, 32, 39, 42, 43, 45, 47/ <i>Lr</i> 2b, 2c, 3, 10, 11, 12, 13, 14a, 14b, 14ab, 16, 17a, 17b, 18, 20, 21, 22a, 22b, 27, 30, 33, 34, 35, 36, 37, 38, 40, 44, 48, 49
12-5	<i>Lr</i> 1, 2a, 9, 10, 13, 15, 19, 24, 25, 28, 29, 32, 36, 39, 42, 43, 45, 47/ <i>Lr</i> 2b, 2c, 3, 11, 12, 14a, 14b, 14ab, 16, 17a, 17b, 18, 20, 21, 22a, 22b, 23, 26, 27+31, 30, 33, 34, 35, 37, 38, 40, 44, 46, 48,49
12-6	<i>Lr</i> 1, 2a, 9, 10, 11, 14b, 14ab, 15, 17a, 17b, 18, 19, 23, 24, 25, 27+31, 28, 29, 32, 36, 38, 39, 42, 43, 45, 47/ <i>Lr</i> 2b, 2c, 3, 12, 13, 14a, 16, 20, 21, 22a, 22b, 30, 33, 34, 35, 37, 40, 44, 46, 48, 49
12-7	<i>Lr</i> 1, 2a, 9, 13, 15, 19, 24, 25, 28, 29, 32, 36, 39, 42, 43, 45, 47/ <i>Lr</i> 2b, 2c, 3, 10, 11, 12, 14a, 14b, 14ab, 16, 17a, 17b, 18, 20, 21, 22a, 22b, 23, 26, 27+31, 30, 33, 34, 35, 37, 38, 40, 44, 46, 48, 49
12-8	<i>Lr</i> 1, 2a, 9, 10, 13, 19, 23, 24, 25, 28, 29, 32, 36, 39, 42, 43, 45, 47/ <i>Lr</i> 2b, 2c, 3, 11, 12, 14a, 14b, 14ab, 15, 16, 17a, 17b, 18, 20, 21, 22a, 22b, 26, 27+31, 30, 33, 34, 35, 37, 38, 40, 44, 46, 48, 49
16	<i>Lr</i> 1, 2a, 2b, 2c, 3, 9, 10, 11, 13, 14, 14b, 14ab, 15, 16, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 32, 34, 36, 37, 38, 39, 42, 44, 45, 47, 48/ <i>Lr</i> 12, 22a, 22b, 33, 35, 49
16-1	<i>Lr</i> 1, 2a, 2b,3, 9,10,11,13,14b,14ab,15,16,17,19,21,24,25,26,27,29,30,32,34,36, 37, 38, 39, 42, 44, 47, 48/ <i>Lr</i> 2c, 12, 14a, 18, 20, 22a, 22b, 23, 33, 35, 49
17	<i>Lr</i> 3, 9, 10, 19, 23, 24, 25, 26, 27+31, 28, 29, 36, 42, 43, 45, 47 / <i>Lr</i> 1, 11, 12, 13, 14a, 15, 16, 17a, 17b, 18, 20, 21, 22a, 22b, 34
20	<i>Lr</i> 3, 9, 10, 15, 16, 17a, 19, 23, 24, 25, 26, 27+31, 28, 29, 36, 42, 43, 45, 47/ <i>Lr</i> 1, 2a, 2b, 2c, 11, 12, 13, 14a, 17b, 18, 20, 21, 22a, 22b, 33, 34, 37, 38, 40, 44, 48, 49
63	<i>Lr</i> 1, 2a, 2b, 2c, 3, 9, 10, 12, 13, 14a, 14b, 14ab, 15, 16, 17a, 17b, 18, 19, 21, 22a, 22b, 23, 24, 25, 26, 28, 29, 30, 32, 33, 34, 35, 36,37, 38, 39, 40, 42, 43, 44, 45, 47, 48, 49/ <i>Lr</i> 11, 20, 27+31, 35
77	<i>Lr</i> 9, 10, 19, 23, 24, 25, 26, 27+31, 28, 29, 32, 36, 39, 42, 43, 45/ <i>Lr</i> 1, 2a, 2b, 2c, 3, 10, 11, 12, 13, 14a, 14b, 14ab, 15, 16, 17, 18, 20, 21, 22a, 22b, 30, 33, 35, 37, 38, 44, 48, 49
77-1	<i>Lr</i> 9, 17,17a, 17b, 19, 23, 24, 25, 27+31, 28, 29, 32, 36, 39, 42, 43, 45, 47/ <i>Lr</i> 1, 2a, 2b, 2c, 3, 10, 11, 12, 13, 14a, 14b, 14ab, 15, 16, 18, 20, 21, 22a, 22b, 26, 30, 33, 35, 37, 38, 44, 48, 49
77-2	<i>Lr</i> 9, 19, 24, 25, 26, 28, 29, 32, 36, 39, 42, 43, 44, 45, 47/ <i>Lr</i> 1,2a, 2b, 2c, 3,10, 11, 12,13, 14a, 14b, 14ab, 15, 16, 17a, 17b, 18, 20, 21, 22a, 22b, 23, 27+31, 30, 33, 34, 35, 37, 38, 40, 48,49
77-3	<i>Lr</i> 9,19,20,23,24, 25, 27+31, 28, 29,32, 36, 39, 42, 43,45, 47/ <i>Lr</i> 1,2a,2b, 2c,3, 10, 11, 12,13, 14a, 14b, 14ab,15, 16, 17, 17a, 17b, 18, 21,22a, 22b,26, 30, 33, 35, 37, 38, 44, 48, 49
77-4	<i>Lr</i> 9, 19, 20, 24, 25, 26, 28, 29, 32, 36, 39, 45, 47/ <i>Lr</i> 1, 2a, 2b, 2c, 3, 10, 11, 12, 13, 14a, 14b, 14ab, 15, 16, 17, 18, 21, 22a, 22b, 23, 27+31, 30, 33, 34, 35, 36, 37, 38, 40, 44, 48, 49
77-5	<i>Lr</i> 9, 19, 24, 25, 28, 29, 32, 39, 42, 43, 45, 47/ <i>Lr</i> 1,2a, 2b, 2c, 3,10, 11, 12,13, 14a, 14b, 14ab, 15, 16, 17a, 17b, 18, 20, 21, 22a, 22b, 23, 26, 27, 30, 33, 34, 35, 36, 37, 38, 40, 44,48, 49
77-6	<i>Lr</i> 9, 18, 19, 20, 24, 25, 28, 29, 32, 39, 40, 42, 45, 47/ <i>Lr</i> 1, 2a, 2b, 2c, 3, 10, 11, 12, 13, 14a, 14b, 14ab, 15, 16, 17a, 21, 22a, 22b, 23, 26, 27+31, 30, 33, 34, 35, 36, 37, 38, 43, 44, 48, 49
77-7	<i>Lr</i> 18, 19, 24, 25, 28, 29, 32, 39, 40, 42, 45, 47/ <i>Lr</i> 1, 2a, 2b, 2c, 3, 9, 10, 11, 12, 13, 14a, 14b,

	<i>14ab, 15, 16, 17a, 17b, 20, 21, 22a, 22b, 23, 26, 27+31, 30, 33, 34, 35, 36, 37, 38, 43, 44, 48, 49</i>
77-8	<i>Lr9, 23, 24, 25, 26, 27+31, 28, 29, 32, 36, 39, 45, 47/Lr1 2a, 2b, 2c, 3a, 10, 11, 13, 14a, 14b, 14ab, 15, 16, 17, 18, 19, 20, 21, 22a, 22b, 30, 33, 35, 37, 38, 44, 48, 49</i>
77-9	<i>Lr2a, 2b, 2c, 9, 19, 24, 25, 28, 32, 39, 42, 45, 47/Lr1, 3, 10, 11, 12, 13, 14a, 14b, 14ab, 15, 16, 17a, 17b, 18, 20, 21, 22a, 22b, 23, 26, 27+31, 30, 33, 34, 35, 36, 37, 38, 44, 46, 48, 49</i>
77-10	<i>Lr2a, 2b, 2c, 9, 19, 24, 25, 32, 39, 42, 45, 47/Lr1, 3, 10, 11, 12, 13, 14a, 14b, 14ab, 15, 16, 17a, 17b, 18, 20, 21, 22a, 22b, 23, 26, 28, 27+31, 30, 33, 34, 35, 36, 37, 38, 44, 46, 48, 49</i>
77-11	<i>Lr2a, 2b, 2c, 9, 19, 24, 25, 26, 28, 29, 32, 36, 39, 42, 45, 47, 57/Lr1, 3, 10, 11, 12, 13, 14a, 14b, 14ab, 15, 16, 17a, 17b, 18, 20, 21, 22a, 22b, 23, 27+31, 30, 33, 34, 35, 37, 38, 44, 46, 48, 49, 51</i>
77-12	<i>Lr2a, 2b, 2c, 9, 19, 20, 24, 25, 28, 32, 39, 42, 45, 47/Lr1, 3, 10, 11, 12, 13, 14a, 14b, 14ab, 15, 16, 17a, 17b, 18, 20, 21, 22a, 22b, 23, 26, 27+31, 30, 33, 34, 35, 36, 37, 38, 44, 46, 48, 49</i>
77-A	<i>Lr9, 17, 19, 23, 24, 25, 26, 27+31, 28, 29, 32, 36, 39, 42, 43, 45, 47/Lr1, 2a, 2b, 2c, 3, 10, 11, 12, 13, 14a, 14b, 14ab, 15, 16, 18, 20, 21, 22a, 22b, 30, 33, 35, 37, 38, 44, 48, 49</i>
77A-1	<i>Lr9, 17, 19, 20, 23, 24, 25, 26, 27+31, 28, 29, 32, 36, 39, 42, 43, 45, 47/Lr1, 2a, 2b, 2c, 3, 10, 11, 12, 13, 14a, 14b, 14ab, 15, 16, 18, 21, 22a, 22b, 30, 33, 35, 37, 38, 44, 48, 49</i>
104	<i>Lr2a, 9, 10, 13, 15, 19, 20, 23, 24, 25, 26, 27, 28, 29, 32, 39, 42, 43, 45, 47/Lr1, 2b, 2c, 3, 11, 12, 14a, 14b, 14ab, 16, 17a, 17b, 18, 21, 22a, 22b, 30, 33, 34, 35, 36, 37, 38, 40, 44, 48, 49</i>
104-1	<i>Lr2a, 9, 15, 19, 24, 25, 26, 28, 29, 32, 39, 42, 43, 45, 47/Lr1, 2b, 2c, 3, 10, 11, 12, 13, 14a, 14b, 14ab, 16, 17a, 17b, 18, 20, 21, 22a, 22b, 23, 27, 30, 33, 34, 35, 36, 37, 38, 40, 44, 48, 49</i>
104-2	<i>Lr9, 10, 13, 15, 19, 20, 24, 25, 28, 29, 32, 36, 39, 42, 43, 45, 47/Lr1, 2a, 2b, 2c, 3, 11, 12, 14a, 14b, 14ab, 16, 17a, 17b, 18, 21, 22a, 22b, 23, 26, 27+31, 30, 33, 34, 35, 37, 38, 40, 44, 48, 49</i>
104-3	<i>Lr9, 10, 13, 15, 17a, 19, 24, 25, 27+31, 28, 29, 32, 36, 39, 42, 43, 45, 47/Lr1, 2a, 2b, 2c, 3, 11, 12, 14a, 14b, 14ab, 16, 17b, 18, 20, 21, 22a, 22b, 23, 26, 30, 33, 34, 35, 37, 38, 40, 44, 48, 49</i>
104-4	<i>Lr2a, 3, 9, 15, 19, 24, 25, 28, 32, 39, 42, 43, 45, 47/Lr1, 2b, 2c, 10, 11, 12, 13, 14a, 14b, 14ab, 16, 17a, 17b, 18, 20, 21, 22a, 22b, 23, 26, 27+31, 29, 30, 33, 34, 35, 36, 37, 38, 40, 44, 46, 48, 49, 51, 57</i>
104A	<i>Lr2a, 9, 10, 15, 19, 21, 23, 24, 25, 26, 27, 28, 29, 32, 39, 42, 43, 45, 47/Lr1, 2b, 2c, 3, 11, 12, 13, 14a, 14b, 14ab, 16, 17a, 17b, 18, 20, 22a, 22b, 30, 33, 34, 35, 36, 37, 38, 39, 40, 44, 48, 49</i>
104B	<i>Lr2a, 9, 15, 19, 20, 24, 25, 26, 28, 29, 32, 39, 42, 43, 45, 47/Lr1, 2b, 2c, 3, 10, 11, 12, 13, 14a, 14b, 14ab, 16, 17a, 17b, 18, 21, 22a, 22b, 23, 27, 30, 33, 34, 35, 36, 37, 38, 40, 44, 48, 49</i>
106	<i>Lr1, 2a, 2b, 3, 9, 10, 11, 12, 13, 14a, 14b, 14ab, 15, 16, 17, 18, 19, 21, 22a, 22b, 23, 24, 25, 26, 27+31, 30, 33, 37, 38, 39, 44, 47, 48/Lr2c, 20, 35</i>
107	<i>Lr1, 3, 9, 10, 19, 20, 23, 24, 25, 26, 27+31, 28, 29, 30, 32, 36, 38, 39, 40, 42, 43, 45, 46, 47/Lr2a, 2b, 2c, 11, 12, 13, 14a, 18, 15, 21, 22a, 22b, 33, 34, 35, 37, 38, 40</i>
107-1	<i>Lr1, 3, 9, 10, 19, 20, 23, 24, 25, 27+31, 28, 29, 30, 32, 36, 38, 39, 40, 42, 43, 45, 46, 47/Lr2a, 2b, 2c, 11, 12, 13, 14a, 18, 15, 21, 22a, 22b, 26, 33, 34, 35, 37, 38, 40</i>
108	<i>Lr3, 9, 10, 15, 17a, 17b, 19, 23, 24, 25, 26, 27+31, 28, 29, 32, 47/Lr1, 2a, 2b, 2c, 11, 12, 13, 14a, 16, 18, 20, 21, 22a, 22b, 33, 34, 35, 36, 37, 38, 44, 48, 49</i>
108-1	<i>Lr3, 9, 10, 17a, 17b, 19, 23, 24, 25, 26, 27+31, 29, 39, 42, 45, 47/Lr1, 2a, 2b, 2c, 11, 12, 13, 14a, 15, 16, 18, 20, 21, 22a, 22b, 33, 34, 35, 36, 37, 38, 44, 48, 49</i>
162	<i>Lr1, 9, 13, 14ab, 15, 19, 20, 21, 23, 24, 25, 26, 27+31, 28, 29, 32, 36, 38, 39, 40, 42, 43, 45, 46, 47/Lr2a, 2b, 2c, 3, 10, 11, 12, 14a, 14b, 16, 17a, 17b, 18, 22a, 22b, 30, 33, 34, 35, 37, 44, 48, 49</i>
162-1	<i>Lr1, 9, 13, 14ab, 15, 19, 21, 23, 24, 25, 27+31, 28, 29, 32, 36, 38, 39, 40, 42, 43, 45, 46, 47/Lr2a, 2b, 2c, 3, 10, 11, 12, 14a, 14b, 16, 17a, 17b, 18, 20, 22a, 22b, 26, 30, 33, 34, 35, 37, 44, 48, 49</i>
162-2	<i>Lr1, 9, 13, 14ab, 15, 19, 20, 21, 23, 24, 25, 27+31, 28, 29, 32, 36, 38, 39, 40, 42, 43, 45, 46, 47/Lr2a, 2b, 2c, 3, 10, 11, 12, 14a, 14b, 16, 17a, 17b, 18, 22a, 22b, 26, 30, 33, 34, 35, 37, 44, 48, 49</i>
162-3	<i>Lr1, 9, 10, 13, 14ab, 15, 19, 20, 21, 23, 24, 25, 26, 27+31, 28, 29, 32, 36, 38, 39, 40, 42, 43, 45, 46, 47/Lr2a, 2b, 2c, 3, 11, 12, 14a, 14b, 16, 17a, 17b, 18, 22a, 22b, 30, 33, 34, 35, 37, 44, 48, 49</i>
162A	<i>Lr1, 9, 13, 14ab, 15, 19, 21, 23, 24, 25, 26, 27+31, 28, 29, 32, 36, 38, 39, 40, 42, 43, 45, 46, 47/Lr2a, 2b, 2c, 3, 10, 11, 12, 14a, 14b, 16, 17a, 17b, 18, 20, 22a, 22b, 30, 33, 34, 35, 37, 44, 48, 49</i>

**Table 5** Analysis of divergence of genetic variation in populations of *Puccinia triticina*

Population	Number of samples collected	Pathotypes identified	$N_a \pm SD$	$N_e \pm SD$	$h \pm SD$	$I \pm SD$	$P_p$ (%)
Madhya Pradesh	238	03	1.61±0.49	1.43±0.39	0.24±0.20	0.36±0.29	61.54
Gujarat	253	02	1.41±0.49	1.29±0.35	0.17±0.20	0.24±0.30	41.03
Haryana	138	04	1.44±0.50	1.34±0.42	0.18±0.22	0.27±0.31	43.59
Karnataka	437	10	1.85±0.36	1.55±0.38	0.31±0.18	0.46±0.25	84.62
Bihar	246	03	1.28±0.45	1.24±0.40	0.12±0.21	0.18±0.29	28.21
Uttarakhand	155	02	1.18±0.38	1.12±0.27	0.07±0.16	0.11±0.23	17.95
Himachal Pradesh	315	04	1.44±0.50	1.24±0.33	0.15±0.18	0.22±0.26	43.59
Tamil Nadu	419	12	1.82±0.39	1.56±0.35	0.32±0.17	0.47±0.24	82.05
Nepal	148	02	1.28±0.45	1.20±0.32	0.11±0.18	0.17±0.27	28.21
Maharashtra	362	02	1.25±0.44	1.8±0.31	0.16±0.18	0.15±0.26	25.64
Punjab	145	02	1.20±0.40	1.15±0.28	0.08±0.16	0.12±0.25	20.51
Uttar Pradesh	124	02	1.26±0.44	1.18±0.31	0.11±0.18	0.15±0.26	25.64
Total	2980	48	1.97±0.16	1.59±0.33	0.34±0.15	0.50±0.19	97.44

$N_a$  = Observed number of alleles; SD = Standard deviation;  $N_e$  = Effective number of alleles;  $h$  = Nei's gene diversity;  $I$  = Shannon's Information index;  $P_p$  = percentage of polymorphic loci



**Table 6** Analysis of molecular variance (AMOVA) among and within the populations of *Puccinia triticina* pathotypes

Source of variation	Degree of freedom	Observed partition*	
		Variance	Variation (%)
Among populations	11	0.860	12.57
Within Populations	36	5.985	87.43
Total	47	6.846	100
*P value = 0.05			


**Figure legends:**

**Figure 1.** Virulence frequency (%) of leaf rust pathotypes on Indian leaf rust differentials.


**Figure 2.** Dendrogram of *Puccinia triticina* pathotypes based on their virulence on 25 leaf rust differentials. Alphabets at the base of the branch indicate the name of the major group.

**Figure 3.** Dendrogram of *Puccinia triticina* pathotypes based on molecular marker data. Alphabets at the base of the branch indicate the name of the major group.


**Figure 4.** Structure analysis of 48 South Asian *Pt* pathotypes. Pathotypes indicated in green and red colours indicate two different allelic patterns among the pathotypes. Pathotypes sharing more than 70% allelic frequency in each category were grouped into a particular subpopulation group and those sharing less than 70% were considered as admixture population.




**Figure 1**



**Figure 2**



**Figure 3**



**Figure 4**

**Supplementary files:**

**Supplementary file** Name of the *Puccinia triticina* pathotypes based on Indian leaf rust differentials and Grouping of *Pt* pathotypes based on allelic frequency through Structure programme.

**Supplementary figure** Allelic pattern among *Puccinia triticina* pathotypes generated with SSR marker, SSR TATTG-60